



International
Animal Health Products
THE AUSTRALIAN COMPANY

BUFFERED IODINE™ SPRAY

AN AID IN THE CONTROL AND TREATMENT OF TOPICAL, BACTERIAL AND FUNGAL INFECTIONS IN ANIMALS.

PRESENTATION: Dark yellow liquid.

ACTIVE CONSTITUENTS: Each L contains:
POVIDONE IODINE 50 g/L equivalent to 5 g/L available iodine

PROPERTIES:

About 70% of cuts and wounds occur on the lower limb of animals, which usually heal considerably slower than skin wounds on the upper body, often delayed by contamination and infection. Lacerations, wounds, skin abrasions and low grade bacterial and fungal infections of the skin are the most common problems of domestic and other animals.

FARNAM® BUFFERED IODINE™ SPRAY is a safe and effective topical spray suitable for general antiseptic purposes on the skin of the most animals. It can be used to control topical infection of the skin, keeping wounds dry and covered with a protective residue film of iodine to provide prolonged activity. **BUFFERED IODINE SPRAY** may be used in numerous surgical procedures and as a topical agent for various dermal and mucosal infections such as wounds, cuts, abrasions, burns, ringworm, navel disinfection in newly born animals, castration and docking wounds, udder wounds and cut teats.

BUFFERED IODINE SPRAY contains “safe” povidone iodine that does not irritate skin or wound surface if used as recommended. **BUFFERED IODINE SPRAY** in the quiet operating pump action pack is ideal for spray on the horses and other animals that often become frightened by the ‘hissing’ sound of pressure-pack topical sprays.

DIRECTIONS FOR USE:

DO NOT APPLY TO EYES OR MUCOUS MEMBRANES.

Government regulations limit iodine residues in milk. Thoroughly wash residues from teats and udder wounds before milking.

Shake bottle and hold upright. Apply **BUFFERED IODINE SPRAY** using the economic pump action. Hold bottle about 15 cm from area to be treated. Spray the affected area twice daily until healed. Avoid application of iodine basic antiseptics to the skin of cats as ingestion from licking a wound may result in allergic reaction to iodine.

MEAT WITHHOLDING PERIOD: NIL.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): ESI not required.

POISONS SCHEDULE: NOT APPLICABLE.

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY.

PACK SIZES: 500 mL spray pack.