

Fidos Closasole 10kg Tablets

COMPOSITION: Levamisole (as hydrochloride) 42.2mg, Niclosamide 1000mg.

ACTIONS: The mechanism of action of Niclosamide appears to be by the uncoupling of oxidative phosphorylation of the cellular electron transport system and by inhibition of ATP synthesis in the parasitic worm. Levamisole acts as a cholinergic agonist and paralyzes the worm allowing its expulsion; however, there is evidence of more than one mode of action. Niclosamide is poorly absorbed from the gut. Levamisole is absorbed from the gut and is metabolised by the liver.

INDICATIONS: Roundworm (*Toxocara* & *Toxascaris* sp.), Hookworm (*Uncinaria* and *Ancylostoma* sp.), and Tapeworm (*Dipylidium* and *Taenia* sp.), in dogs and cats.

CONTRAINDICATIONS: Known sensitivity to levamisole or niclosamide.

PRECAUTIONS: Discontinue use if unsteady gait or depressed appetite occurs after administration. Use with caution in debilitated animals and animals with reduced hepatic function. To prevent vomiting, administer with food and divide the dose.

FIRST AID: If poisoning occurs, contact a doctor or a Poisons Information Centre. Phone Australia 131 126.

DISPOSAL: Dispose of empty container by wrapping in paper and placing in garbage.

DOSAGE AND ADMINISTRATION: Administer with a light meal or crushed in food.

Adult dogs and cats: 1 tablet /10Kg bodyweight once. Repeat every three months if necessary throughout life.

Bitches and Queens: 1 tablet /10Kg bodyweight, two treatments at an interval of 14 days during gestation and lactation.

PRESENTATION: Tablets: 10's, 100's.

STORAGE: Store below 30°C (Room temperature). Protect from light. Replace lid tightly after use.

POISONS SCHEDULE: S6

APVMA NUMBER: 37422