

Frontline Plus

FRONTLINE PLUS is indicated for the treatment and prevention of flea infestation and control of brown dog ticks, paralysis ticks and biting lice on dogs and puppies.

FRONTLINE PLUS has a rapid onset of action and kills re-infestations with newly acquired adult fleas for at least one month. FRONTLINE PLUS also prevents the development of flea eggs, larvae and pupae produced by adult fleas acquired for three months after treatment.

FRONTLINE PLUS controls brown dog ticks for up to one month, and paralysis ticks for up to two weeks after application. Daily searching of dogs is necessary to minimise the risk of tick paralysis. Searching should continue throughout the paralysis tick season.

FRONTLINE PLUS may be used in the treatment and control of Flea Allergy Dermatitis in dogs.

FRONTLINE PLUS may be used as the only product for flea control as it breaks the life cycle on the animal and in its surroundings by killing all stages of fleas.

Dogs may be treated with FRONTLINE PLUS following bathing and shampooing once they are dry.

The product will spread over the coat of the animal within 24 hours. Once spread, FRONTLINE PLUS is water fast. It is recommended not to bath or shampoo dogs during the 48 hours after treatment. Thereafter, bathing, water immersion or exposure to sunlight will not reduce the efficacy of FRONTLINE PLUS.

Safety:

FRONTLINE PLUS has demonstrated a significant margin of safety for use in dogs and puppies over 8 weeks of age and in breeding, pregnant or lactating bitches.

Treatment Frequency:

Fleas: Monthly application is recommended. All animals in a household should be treated at the same time (for household cats use FRONTLINE PLUS for Cats). It is especially important to treat all household animals if any pet is allergic to fleas (flea allergy dermatitis).

Paralysis Tick: Two weekly applications are recommended. While FRONTLINE PLUS provides control of paralysis tick for up to 2 weeks, use of this

product does not guarantee prevention of tick paralysis because ticks are not killed immediately after attachment. Daily searching of dogs is necessary to minimise the risk of tick paralysis. See your veterinarian for advise on searching techniques and tick seasonality.

Brown Dog Tick: Monthly application is recommended.

Biting Lice: Treat all household dogs at the same time.



Frontline Plus - Spray

For the treatment and prevention of flea and tick (including Ixodes holocyclus) infestations on cats and dogs.

Active Ingredient: 2.5 g/L Fipronil

APPLICATION METHOD: Ensure complete coverage to skin level. Massage the coat all over to make sure the spray gets right down to the skin.

DOSE RATE: Fleas: 3-6 mL/kg = 2-4 spray pumps/kg
Flea Allergy Dermatitis: 6 mL/kg = 4 spray pumps/kg
Paralysis Ticks: 6 mL/kg = 4 spray pumps/kg

PACK SIZES: Available in 3 pack sizes: 100mL, 250mL and 500mL.

