



Canimax offers an all-in-one solution for the treatment of all major worms that can harm dogs including heartworm prevention. Canimax is a palatable tablet that can be easily administered every 6 weeks, Canimax will provide an effective and safe solution to the main intestinal parasite problems that dog can be faced with.

# CANIMAX<sup>®</sup>

## Palatable Broad Spectrum Wormer



### WHAT IS CANIMAX?

- ✓ Prevents Heartworm
- ✓ Treats Roundworm
- ✓ Treats Hookworm
- ✓ Treats Whipworm
- ✓ Treats Tapeworm (including hydatids)
- ✓ Liver Flavoured chewable tablet

### WHY CHOOSE CANIMAX?

#### Broad spectrum specialist worming product

CANIMAX prevents heartworm and kills all intestinal worms (roundworm, hookworm, whipworm and tapeworm). One easy dose will protect your dog for 6 weeks from all worms that can be dangerous to your pet. Canimax is safe to use in puppies from 6 weeks of age, lactating and pregnant bitches, older dogs and collie dogs. This means peace of mind that your pet is being treated safely without the risk of complications.

#### Available in 3 sizes for dogs

Small (0-5kg), Medium (5-10kg) and Large (10-20kg)

#### Liver flavoured treat

CANIMAX is a tasty liver flavour chewable tablet. It converts worming into a treat. Tablets are quarter scored to allow for flexible dosing in multi-pet households.

### WHEN TO USE CANIMAX?

Canimax can be used from 6 weeks of age and simply retreat every 6 weeks

- SMALL DOGS: 1 TABLET FOR 5KG BW
- MEDIUM DOGS: 1 TABLETS FOR 10KG BW
- LARGE DOGS: 1 TABLET FOR 20KG BW

### WHY DOES MY DOG NEED TO BE WORMED?

Heartworm disease is a serious and potentially fatal condition. It can go undetected for many years and by the time it is diagnosed the worms may have caused significant damage to the heart and lungs.

The disease is carried from one infected dog to healthy dogs by mosquitos and both indoor and outdoor dogs on mainland Australia are at risk.

Wherever you live, intestinal worms (roundworm, hookworm, whipworm and tapeworm) can cause illness in pets and potentially humans.

Dogs can pick up intestinal worms by transmission from the mother during pregnancy or following birth, by ingestion from contaminated food, from faeces, fleas, skin penetration, or from close contact with another animal.

These worms can cause diarrhoea, ill thrift a pot-bellied appearance, failure to thrive, and a dry shaggy coat in dogs.

Some intestinal worms affecting dogs can potentially be transmitted to people. Children who play in soil contaminated with dog faeces are at risk of accidental ingestion of worm eggs.

Treating your dog with Canimax is a simple solution to these problems.

**Virbac**  
ANIMAL HEALTH

*Passionate about animal health*

### Palatable Broad Spectrum Wormer

#### INDICATIONS

- Treatment of Roundworm (*Toxascaris leonina*, *Toxocara canis*), Hookworm (*Ancylostoma caninum*, *Uncinaria stenocephala*), Whipworm (*Trichuris* spp.) and Tapeworm (*Dipylidium caninum*, *Taenia* spp.) including hydatids (*Echinococcus granulosus*) in dogs. Prevention of heartworm (*Dirofilaria immitis*).

#### RECOMMENDED FOR USE IN

- All dogs from 6 weeks of age

#### PRESENTATION

Available in 3 sizes for dogs:

- Small Dog (one tablet per 0-5kg body weight)
- Medium Dog (one tablet per 5 – 10kg body weight)
- Large Dog (one tablet per 20kg body weight)

The above presentations are in 4 tablet packs.

#### COMPOSITION

	Small Dog	Medium Dog	Large Dog
Abermectin	0.05mg	0.1 mg	0.2mg
Oxibendazole	112.5mg	225mg	450mg
Praziquantel	25mg	50mg	100mg

#### DOSAGE & DIRECTIONS

You must consult a veterinarian before starting any heartworm prevention as your dog might require testing to establish whether or not heartworm is already present. If the pet is already being treated with an existing heartworm prevention product it is safe to switch to Canimax when the next dose is due.

- PUPPIES: Commence treatment at 6 weeks of age and retreat every 6 weeks.
- ADULTS: Retreatment is required every 6 weeks.
- Dogs should not be fed, or allowed to feed on, offal from any species.
- Dogs in hydatid areas should be treated every 6 weeks. Where whipworm is a problem treat every 6-8 weeks after 3 months of age.

