



Kills and controls ALL intestinal worms Endogard is for the treatment and prevention of roundworm, hookworm, whipworm and tapeworm, including hydatids. Endogard is safe to use with any heartwormer for dogs on the market.* It is an integral part of an effective parasite control program for dogs. Tasty Liver Flavour dogs will love the liver flavoured chewable tablet so administration is easy.

endogard

Intestinal Wormer



WHAT IS ENDOGARD?

- ✓ Broad spectrum allwormer
- ✓ Tasty liver flavour
- ✓ Easy to administer
- ✓ Safe

WHY CHOOSE ENDOGARD?

Kills and controls all intestinal worms

Endogard is for the treatment and prevention of roundworm, hookworm, whipworm and tapeworm, including hydatids. Endogard is safe to use with any heartwormer for dogs on the market. It is an integral part of an effective parasite control program for dogs

Tasty liver flavour

Dogs will love the liver flavoured chewable tablet so administration is easy.

Flexible dosing

Endogard chewable tablets are quarter scored to allow for flexible dosing in multi-pet households.

WHY DOES MY DOG NEED TO BE WORMED?

Wherever you live in Australia, intestinal worms (roundworm, hookworm, whipworm and tapeworm) can cause illness in pets and potentially humans.

Dogs can pick up intestinal worms by transmission from the mother during pregnancy or following birth, by ingestion from contaminated food, from faeces, fleas, skin penetration, or from close contact with another animal.

These worms can cause diarrhoea, ill thrift, a pot-bellied appearance, failure to thrive, and a dry shaggy coat in dogs.

Some intestinal worms affecting dogs can potentially be transmitted to people. Children who play in soil contaminated with dog faeces are at risk of accidental ingestion of worm eggs.

By treating your dog with Endogard Palatable Allwormer for Dogs you can prevent these parasites from harming you, your family and your pet.

SAFETY OF ENDOGARD WITH COMMONLY USED INSECTICIDES

Monash University conducted a confirmatory safety study with Virbac Endogard Palatable Allwormer when administered concurrently with insecticides*.

Method

A total of 16 pups were used (9 greyhounds and 7 crossbred pups) ranging in age from 3-7 months. These pups were divided into 3 groups, 2 of which were given a combination of commonly used insecticides as outlined below, as well as the test material. The third group was only given Virbac Endogard Palatable Allwormer. The group of actives in group 2 covered the routine chemicals used as dog ectoparasiticides. The insecticides used were applied as per manufacturer's recommendations.

Timetable for insecticide and Endogard administration				
	Day 0	Day 14	Day 28	Day 35
Group 1	HeartGuard® Program® Frontline® & Endogard	Endogard	Program® Frontline® Proban® & Endogard	Final Check
Group 2	Dimmitrol® Preventic® Care® & Endogard	Endogard	Care® & Endogard	Final Check
Group 3	Endogard	Endogard	Endogard	Final Check

All dogs were observed daily during the trial. They were also individually examined at 0, 6, 24 and 48 hours after each administration.

Results

No abnormal developments were noted at any time point, examination or observation.

Conclusion

This study confirms that there was no evidence of deleterious effects on the pups. Virbac Endogard Palatable Allwormer is safe to use in dogs under most conditions, particularly as the trial conditions would normally exceed routine use of insecticides, heartworm treatment and gastrointestinal worming treatment.



Passionate about animal health

INDICATIONS

- Treatment and control of roundworm (*Toxascaris leonina*, *Toxacara canis*), hookworm (*Ancylostoma caninum*, *Ucinaria stenocephala*), whipworm (*Trichuris spp*) and tapeworm (*Dipylidium caninum*, *Taenia spp*), including hydatids (*Echinococcus granulosus*) in dogs. Does not control heartworm.

RECOMMENDED FOR USE IN

- All dogs from 6 weeks of age

PRESENTATION

Small dog (0-5kg) tablet
(25mg praziquantel and 112.5mg oxibendazole)

Medium dog (5-10kg) tablet
(50mg praziquantel and 225mg oxibendazole)

Large dog (10-20kg) tablet
(100mg praziquantel and 550mg oxibendazole)

Available in cartons and bulk packs

COMPOSITION

Contains:

Praziquantel	5mg/kg
Oxibendazole	22.5mg/kg

DIRECTIONS FOR USE

Young Puppies:

Treat according to weight at 2, 4, 8 and 12 weeks of age.

Adult Dogs:

Treat according to weight every three months. If your pet is a breeding bitch, or located in a known hydatid area or if whipworm is a problem, more frequent administration is required.

